

1. (32 pts) The position function of a particle is given by  $s(t) = 4t^2$  on the interval  $1 \leq t \leq 16$ , where position is in meters and time is in seconds.

(a) Determine the particle's velocity function  $v(t)$ . Include the correct unit of measurement.

**Solution:**

$$s(t) = 4t^2 \quad t = 4t^{1=2} \quad t$$

$$v(t) = s'(t) = 2t^{1=2} \quad 1 = \boxed{\frac{2}{t} \quad 1 \text{ m/s}}, \quad 1 < t < 16$$

(b) Determine the total distance traveled by the particle on the interval  $1 \leq t \leq 16$ .

- (c) i. Verify that all hypotheses of the Mean Value Theorem are satisfied for the given position function  $s(t) = 4t^2 - t^3$  on the interval  $1 \leq t \leq 16$ .

**Solution:**

$s(t)$  is continuous on  $[1; 16]$  and differentiable on  $(1; 16)$

- ii. Use the Mean Value Theorem to determine all time values  $c$  on the interval  $1 \leq t \leq 16$ , if any, for which the instantaneous velocity of the particle equals the average velocity of the particle on that interval. Include the correct unit of measurement.

**Solution:**

The Mean Value Theorem states that since the hypotheses have been satisfied, there exists at least one number  $c$  on the interval  $(1; 16)$  such that

$$s'(c) = \frac{s(16) - s(1)}{16 - 1} = \frac{0 - 3}{15} = -\frac{1}{5}$$

Therefore, using the result from part (a), we have

$$v(c) = \frac{2}{c} - 1 = -\frac{1}{5}$$

$$\frac{2}{c} = \frac{4}{5}$$

$$c = \frac{5}{2}$$

$$c = \frac{25}{4} \text{ sec}$$

2. (11 pts) Let  $v$  represent a person's walking speed, expressed in miles per hour, and let  $p$  represent the corresponding walking pace, expressed in minutes per mile. The pace can be expressed as the following function of speed:

$$p(v) = \frac{60}{v}; \quad v > 0$$

- (a) Find the linearization  $L(v)$  that approximates  $p(v)$  near  $v = 4$ .

**Solution:**

$$p(v) \quad L(v) = p(4) + p'(4)(v - 4)$$

$$p(4) = \frac{60}{4} = 15$$

$$p'(v) = \frac{60}{v^2} \quad ) \quad p'(4) = \frac{60}{4^2} = \frac{15}{4}$$

$$L(v) = \boxed{15 + \frac{15}{4}(v - 4)}$$

- (b) Use your linearization from part (a) to estimate the walking pace of a person moving at 4.2 miles per hour. Include the correct unit of measurement. You must use linearization to earn credit.

**Solution:**

$$p(4.2) \quad L(4.2) = 15 + \frac{15}{4}(4.2 - 4) = 15 + \frac{15}{4} \cdot \frac{1}{5}$$

3. (26 pts) Consider the function  $f(x) = \sin x + \cos^2 x$  on the interval  $[0; 2\pi]$ .

(a) Identify all critical numbers of  $f$  on the specified interval.

**Solution:**

Critical numbers are values of  $x$  in the domain of  $f$  such that  $f'(x) = 0$  or  $f'(x)$  does not exist. There are no critical numbers of the latter type for this function.

$$f'(x) = \cos x + 2 \cos x (-\sin x) = \cos x(1 - 2 \sin x) = 0$$

$x = \pi/2$  is the only value of  $x$





6. (15 pts) Determine  $g'(x)$  for the function  $g(x) = \sqrt[3]{3x+1}$  by using the definition of derivative. You must obtain  $g'$  by evaluating an appropriate limit to earn credit.

**Solution:**

The definition of derivative indicates that  $g'(x) = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = \lim_{h \rightarrow 0} \frac{\sqrt[3]{3(x+h)+1} - \sqrt[3]{3x+1}}{h}$

7. (22 pts) Consider the function  $h(x) = \frac{\sin x}{x(x-2)}$ .

(a) Find the  $(x; y)$