DISCUSSION PAPERS IN ECONOMICS

Working Paper No. 23-10

!"#$%&'()*$+,-.$'&"*(+,/"&0,1)(+*2)3,,
4$&5*#)+,'(3,!*5"'%,6(&)2'5&"*(+,

!"#$%&'$()*+,")
-&$.(/0$#1),2)3,4,/%5,)6,"45(/)

78#,9(/):;<)=>=:)

Depar men of Economics

Uni ersi of Colorado Bo lder
Bo lder, Colorado 80309

?)78#,9(/):;<)=>=:)!"#$%&'$()*+,")

Simultaneous Equations with Censored Outcomes and Social Interactions

Sutianjie Zhou[▫]

Oct 31, 2023

Abstract

This paper introduces a censored-outcome simultaneous-equation model with social interactions. The construction of the microeconomics foundation for this model is from the equilibrium in a large-network-based game with incomplete information, in which each agent conducts multiple actions and interacts with other agents through a network through a linear quadratic utility function. The su cient condition of the unique Bayesian Nash Equilibrium (BNE) existence is characterized. We also discuss the identi cation of the econometric model. We propose a two-stage method to estimate the model in which we apply the nested pseudo-likelihood (NPL) to estimate the reduced parameters and then derive the structural form parameters by Amemiya Generalized Least Square estimator (AGLS). Monte Carlo simulation shows that the estimation performs well in nite samples. The estimation also shows the feasibility of the computation when the network size is large.

JEL classifcation: C31, C34

Key words: Censored Model, Limited Dependent, Networks, Spatial Autoregressive Models, Simultaneous Equation Models, Social Interaction Models, Tobit model

1 Introduction

In many real economic situations, the outcomes of agents' activities are censored. For example, the spending amount of household annually traveling should be zero or some positive value. Researchers of teenagers' behaviors are interested in how many cigarettes a teenager smokes and how many alcoholic beverages a teenager drinks. Both results are also zeros or positive values. Such result of each agent's certain activity can be in uenced by the same agent's other activities' results and also can be in uenced by other agents' activities'. From the household consumption example, a household's annual traveling spending will in uence its annual entertainment spending and vice

[▫]Department of Economics, University of Colorado at Boulder

versa. And both spending can be impacted by their friends' spending. As for the teenager behavior research example, the number of cigarettes a teenager smokes also impacts how many alcoholic beverages he/she drinks and vice versa. Both the number of cigarettes he/she smokes and the number of alcoholic beverages he/she drinks are in uenced by his/her friends. The multi-action censored outcomes of di erent actions determined by the same agent inter-depend, and agents also impact each other. This paper proposes a simultaneous equation model with censored outcomes and social interactions to characterize the interdependence of an agent's economic decision outcomes both across agents and economic activities.

From the seminal literature, Amemiya (1974) proposed a multivariate Tobit model, which is an extension of Tobin (1958) and Amemiya (1973), and build a simpli ed version of consistent estimation by grouping agents into two types based on the outcomes of there activities. Nelson and Olson (1978) generate a di erent multivariate Tobit model with latent endogenous variables under a more general truncation structure, i.e., not all zeros, and also develop an estimator. The model they propose has one completely observed endogenous variable and one truncated. They derived a reduced-form-based two-stage estimation containing the application of the least square method and likelihood method. Then, Amemiya (1979) extends the simultaneous-equation Tobit model more generally, which combines truncated and non-truncated endogenous outcomes and develops a GLS-based estimator. According to the variance-covariance matrix developing, the estimation process Amemiya (1979) is more e cient than that in Nelson and Olson (1978). Our econometric model is similar to the traditional simultaneous equation model, but the endogenous variables of our structure model are dependent not on the observed outcomes but on the unobserved reservation values.

Our paper is based on the intuition in which each agent conducts various activities at the same time and shows the results, and each agent cannot observe others' outcomes before making decisions, however, their expectation of other agents' outcomes will in uence their behaviors. That means the econometricians and economists need to gure out the Bayesian Nash Equilibrium in the incomplete information network game. Liu (2019) discuss a simultaneous-equation binary model with social interactions under incomplete information and conduct the estimation based on maximum likelihood and nested pseudo-likelihood, which is introduced in Aguirregabiria and Mira (2007). Yang et al. (2018) discuss the incomplete information situation of the single-equation Tobit model with social interactions and compare it to the complete information case, which is an

extension of Xu and Lee (2015), in which a maximum likelihood estimator is developed to estimate the single-equation SAR Tobit model. Other related literature, such as Lee et al. (2014), Lin and Xu (2017), and Yang and Lee (2017) discuss the existence of Bayesian Nash Equilibrium under incomplete information social network in which individuals share rational expectations on others' outcomes.

The rest of this paper is organized as follows. Section 2 clari es the microeconomic foundation of this paper, a network multi-activity game under incomplete information case. Section 3 is the generation of the econometric model of this paper. Section 4 is the estimation. Section 5 is the Monte Carlo simulation. Section 6 concludes.

to write the vector form, we introduce the following notations,

$$\begin{aligned} \mathbf{y}_l &= (y_{1l}; y_{2l}; \ldots; y_{nl})' \\ \mathbf{p}_l &= (p_{1l}; p_{2l}; \ldots; p_{nl})' \\ \mathbf{}_k &= (\ _{1k};\ _{2k}; \ldots;\ _{nk})' \\ \mathbf{}_k &= (\ _{1k};\ _{2k}; \ldots;\ _{nk})' \end{aligned} \tag{5}$$

the vector form can be written as

$$\sum_{l=1} \ _{lk}\mathbf{y}_l = \sum_{l=1} \ _{lk}\mathbf{W}\mathbf{p}_l + \ _k \quad \ _k \tag{6}$$

In the vector form equation (6), the $_{lk}$ represents the interdependence e ect among activities conducted by the same agent. In other words, an agent's underlying intention in activity k will depend on his/her underlying intention in activity l. And obviously, we have $_{kk} = 1$. $_{kk}$ represents the inner-activity's peer e ect, which means an agent's intention of activity k may be impacted by the expected activities' outcomes of the peers in the same activity. $_{kl}$ represents the inter-activity's peer e ect, which means an agent's intention of activity k may be impacted by the expected activities' outcomes of the peers in activity l. Then, we introduce the following notations to write the matrix form

$$\begin{aligned} \mathbf{Y} &= [\mathbf{y}_1; \mathbf{y}_2; \ldots; \mathbf{y}_m] \\ \mathbf{P} &= [\mathbf{p}_1; \mathbf{p}_2; \ldots; \mathbf{p}_m] \\ &= [\ _1;\ _2; \ldots;\ _m] \\ \mathbf{E} &= [\ _1;\ _2; \ldots;\ _m] \end{aligned} \tag{7}$$

then the matrix form can be written as

$$\mathbf{Y} = \mathbf{W}\mathbf{P} + \quad \mathbf{E} \tag{8}$$

If is non-singular, we will have the reduced form of the matrix form of our model, which is

$$\mathbf{Y} = \mathbf{W}\mathbf{P} + \quad \mathbf{E} \tag{9}$$

where $= \ ^{-1}$, $= \ ^{-1}$, and $\mathbf{E} = \mathbf{E}\ ^{-1}$. According to the reduced matrix form of our

model, we can derive the scalar reduced form as

$$y_{ik} = \sum_{l=1}^{\infty} \sum_{j=1}^{\infty} w_{ij} p_{jl} + \quad _{ik} \quad _{ik} \tag{10}$$

where $[\ _{lk}]_{l=1;k=1}^{m;m} =$, $[\ _{ik}]_{i=1;k=1}^{n;m} =$, and $[\ _{ik}]_{i=1;k=1}^{n;m} = E$. Then

$$\begin{aligned} p_{ik} &= E(y_{ik} | \{\$_{jl}\}_{j=1;l=1}^{n;m}) \\ &= E(y_{ik} | \{\$_{jl}\}_{j=1;l=1}^{n;m}; y_{ik} > 0) Pr(y_{ik} > 0 | \{\$_{jl}\}_{j=1;l=1}^{n;m}) \\ &\quad + E(\end{aligned}$$

According to the expression of

Proposition 2.2. If Assumption 1, 2, and 3 holds, then the incomplete information network game has a unique pure strategy BNE, given the equilibrium strategy $y^?$ as

$$y^? = (\quad^0 \quad W)p^? + \tag{18}$$

where the vector of equilibrium beliefs $p^?$ is the unique solution of

$$p = \hbar(p) \tag{19}$$

where

$$\hbar(p) = [\hbar_1(p)^0; \hbar_2(p)^0; \quad ; \hbar_m(p)^0]^0$$

and

$$\hbar_k(p) = [F_k(u_{1k}); F_k(u_{2k}); \quad ; F_k(u_{nk})]^0$$

where

$$u_{ik} = \sum_{l=1}^{\infty} \sum_{lk} \sum_{j \in i} w_{ij} p_{jl} + \quad _{ik}$$

and

$$F_k(u) = u\Phi(\frac{u}{\quad_k}) + \quad _k \ (\frac{u}{\quad_k})$$

where $\Phi(\ )$ and $\ (\ )$ is C.D.F. and P.D.F. of standard normal distribution.

Proof: As $F_k(\ )$ is continuous for all $k \ 2 \ f1; 2; \quad mg$, therefore, $\hbar(\ )$ is continuous, therefore, according to Brouwer xed-point theorem, there at least exist one solution to $p = \hbar(p)$. According to the contraction mapping theorem, the solution to $p = \hbar(p)$ is unique if there exists some kind of norm that the Hessian matrix norm less than one, i.e., $jj@\hbar(p)=@p^0jj < 1$ for some $jj \ \ jj$ then we have

$$\frac{@\hbar(p)}{@p^0} = \begin{bmatrix} \frac{@\hbar_1(\mathbf{p})}{@\mathbf{p}_1^0} & \cdots & \frac{@\hbar_1(\mathbf{p})}{@\mathbf{p}_m^0} \\ \vdots & \ddots & \vdots \\ \frac{@\hbar_m(\mathbf{p})}{@\mathbf{p}_1^0} & & \frac{@\hbar_m(\mathbf{p})}{@\mathbf{p}_m^0} \end{bmatrix}$$

then

$$\begin{aligned}\frac{F_k(u_{ik})}{@p_{jl}} &= {}_{lk}\, w_{ij}\Phi(\frac{u_{ik}}{k}) + \frac{w_{ij}}{k}u_{ik}\ (\frac{u_{ik}}{k}) \quad {}_k w_{ij}\frac{u_{ik}}{k}\frac{1}{k}\ (\frac{u_{ik}}{k})] \\ &= {}_{lk}w_{ij}\Phi(\frac{u_{ik}}{k}) \\ &\leq {}_{lk}w_{ij}\end{aligned}$$

It follows

$$\begin{aligned}\text{jj}\frac{@\hbar(\text{p})}{@\text{p}^0}\text{jj}_1 &\leq \max_{k=1;\cdots m}\ \sum_{l=1}^{\infty}\ \text{j}\ {}_{lk}\text{j}\max_{i=1;\cdots n}\ \sum_{j=1}^{\infty}\text{j}w_{ij}\text{j} = \text{jj}\quad \text{jj}_1\text{jj}\text{W}\text{jj}_1 \\ \text{jj}\frac{@\hbar(\text{p})}{@\text{p}^0}\text{jj}_1 &\leq \max_{l=1;\cdots m}\ \sum_{l=1}^{\infty}\ \text{j}\ {}_{lk}\text{j}\max_{j=1;\cdots n}\ \sum_{i=1}^{\infty}\text{j}w_{ij}\text{j} = \text{jj}\quad \text{jj}_1\text{jj}\text{W}\text{jj}_1\end{aligned}$$

When all the previous assumptions hold, we can derive the uniqueness of the solution from proposition 2.2 according to the completeness and coherency of the model in Tamer (2003). The contraction mapping (19) with a xed point will guarantee the convergence to a consistent estimator in Kasahara and Shimotsu (2012), hence suggesting the Nested Pseudo Likelihood estimation algorithm. We will propose the econometric model based on the incomplete social network game and discuss the identi cation and estimation in the next section.

3 Econometric Model

Suppose $_k = \text{X}\ _k$, where $\text{X} = [\text{x}_1; \text{x}_2; :::; \text{x}_n]^0$ is an $n \quad q$ matrix, representing exogenous variables; and $_k$ is unknown q-dimension parameter vector re ect the direct e ect in the model, then we can consider a simultaneous-equation model with a censored outcome variable $y_{ik} = y_{ik}^{\ } \ 1(y_{ik} > 0)$, where

$$y_{ik} = \sum_{l=1;l\neq k}^{\infty}{}_{lk}y_{il} + \sum_{l=1}^{\infty}{}_{lk}\sum_{j=1;j\neq i}^{\infty}w_{ij}p_{jl} + \text{x}_i\ {}_k \quad {}_{ik}; \tag{20}$$

for $i = 1; \quad ; n$ and $k = 1; \quad ; m$. Let $\text{Y} = [\text{y}_1; \text{y}_2; \quad ; \text{y}_m]$, $\text{P} = [\text{p}_1; \text{p}_2; \quad ; \text{p}_m]$, $\text{X} = [\text{x}_1; \text{x}_2; \quad ; \text{x}_n]^0$, $\text{E} = [\ _1; \ _2; \quad ; \ _m]$, and $W = [w_{ij}]$, where $\text{y}_k = (y_{1k}; y_{2k}; \quad ; y_{nk})^0$, $\text{p}_k = (p_{1k}; p_{2k}; \quad ; p_{nk})^0$, and $_k = (\ _{1k}; \ _{2k}; \quad ; \ _{nk})^0$. Let $\quad = [\ _{kl}]$, an $(m \quad m)$-dimension matrix that re ects the inner e ect cross activities of the same agent, where $_{kk} = 1$ for all k, $\quad = [\ _{kl}]$,

an (m m)-dimension matrix that re ects the peer e ects among agents and activities; and B = [$_1$; ; $_m$], which re ects the direct e ects. In matrix form, Equation (20) can be written as

$$Y = WP + XB E: \tag{21}$$

Based on our Assumption 2, we can extend our assumption that vec(E)jX $N(0;$ $I_n)$. The $(k;l)$-th element of is $_{kl}$ $_k$ $_l$, with $_{kk}$ = 1 for all $k = 1;$; m. If is nonsingular, the reduced form of Equation (21) is

$$Y = WP + XB E ; \tag{22}$$

where = $^{-1}$, B = B $^{-1}$, and E = E $^{-1}$. As vec(E)jX $N(0;$ $I_n)$, we have vec(E)jX $N(0;$ $I_n)$ where = $^{0-1}$ $^{-1}$. The $(k;l)$-th element of is $_{kl}$ $_k$ $_l$, with $_{kk}$ = 1 for all $k = 1;$; m. From equation (22), we can derive the scalar reduced form of the econometric model as

$$y_{ik} = \sum_{l=1}^{m} {}_{lk} \sum_{j=1; j\neq i}^{n} w_{ij} p_{jl} + x_i \; {}_k \; {}_{ik}: \tag{23}$$

Let $d_{ik} = 1$ if $y_{ik} > 0$. Then,

$$\Pr(d_{ik} = 1) = \Pr(y_{ik} > 0) = \Pr(\sum_{l=1}^{m} {}_{lk} \sum_{j=1; j\neq i}^{n} w_{ij} p_{jl} + x_i \; {}_k > {}_{ik}) = \Phi_{ik};$$

where $\Phi_{ik} = \Phi[(\sum_{l=1}^{m} {}_{lk} \sum_{j=1; j\neq i}^{n} w_{ij} p_{jl} + x_i \; {}_k)= {}_k]$. And $\Phi(\,)$ is the cdf of standard normal distribution

$$\begin{aligned} p_{ik} \quad & E(y_{ik}) = E[E(y_{ik} | d_{ik})] \\ = \; & E(y_{ik} | d_{ik} = 1) \Pr(d_{ik} = 1) + E(y_{ik} | d_{ik} = 0) \Pr(d_{ik} = 0) \\ = \; & E(y_{ik} | d_{ik} = 1) \Pr(d_{ik} = 1) \\ = \; & (\sum_{l=1}^{m} {}_{lk} \sum_{j=1; j\neq i}^{n} w_{ij} p_{jl} + x_i \; {}_k)\Phi_{ik} + \; {}_k \; {}_{ik} \end{aligned}$$

where $_{ik}$ = [($\sum_{l=1}^{m}$ $_{lk}$ $\sum_{j=1; j\neq i}^{n}$ $w_{ij} p_{jl}$ + x_i $_k$

are observational equivalent if

$$\left(\sum_{l=1}^{m} e_{lk} \sum_{j=1;j\neq i}^{n} w_{ij}\tilde{p}_{jl} + x_i \tilde{e}_k\right)\tilde{\Phi}_{ik} + e_k \tilde{e}_{ik} = \left(\sum_{l=1}^{m} {}_{lk} \sum_{j=1;j\neq i}^{n} w_{ij} p_{jl} + x_i {}_k\right)\Phi_{ik} + {}_k {}_{ik} \tag{24}$$

for all $i = 1; \quad ; n$ and $k = 1; \quad ; m$. Based on our Assumption 1, 2, and 3, p_{ik} and $\tilde{p}_{ik}$ are xed point solutions to the contraction mapping, which should be identical to each other due to the uniqueness. $p_{ik} = \tilde{p}_{ik}$ for all $i = 1; \quad ; n$ and $k = 1; \quad ; m$. Therefore we have

$$\begin{aligned} &[WP; X][\quad {}^{0}; B\ {}^{0}]^{0} \quad \Phi([WP; X][\quad {}^{0}; B\ {}^{0}]^{0} D_m^{-1}) + \quad ([WP; X][\quad {}^{0}; B\ {}^{0}]^{0} D_m^{-1}) D_m \\ =&[WP; X][e\ {}^{0}; \tilde{B}\ {}^{0}]^{0} \quad \Phi([WP; X][e\ {}^{0}; \tilde{B}\ {}^{0}]^{0} \tilde{D}_m^{-1}) + \quad ([WP; X][e\ {}^{0}; \tilde{B}\ {}^{0}]^{0} \tilde{D}_m^{-1}) \tilde{D}_m \end{aligned} \tag{25}$$

where $\quad$ is the Hadamard (Schur) product of matrices explained in section 7.5 of Horn and Johnson (2012). D_m and $\tilde{D}_m$ are diagonal matrices with diagonal elements $\{ {}_k^2\}_{k=1}^m$, $\{e_k^2\}_{k=1}^m$, and other elements are zeros. Then we can derive the scalar form as

$$\left(\sum_{l=1}^{m} e_{lk} \sum_{j=1;j\neq i}^{n} w_{ij}\tilde{p}_{jl} + x_i \tilde{e}_k\right) + e_k \frac{e_{ik}}{\tilde{\Phi}_{ik}} = \left(\sum_{l=1}^{m} {}_{lk} \sum_{j=1;j\neq i}^{n} w_{ij} p_{jl} + x_i {}_k\right) + {}_k \frac{ {}_{ik}}{\Phi_{ik}} \tag{26}$$

Suppose we denote $L = [\ {}_{ik}/\Phi_{ik}]_{i=1;k=1}^{n;m}$, then if the matrix $[WP; X; L]$ is full column rank, the reduced form parameters $[\quad ; B\ ; D_m]$ are identi able. As

$$[WP; X; L] \ [\quad ; B\ ; D_m] \quad [e\ ; \tilde{B}\ ; \tilde{D}_m] \ = 0 \tag{27}$$

the scalar form results are

$$\begin{aligned} \left(\sum_{l=1}^{m} {}_{lk} \sum_{j=1;j\neq i}^{n} w_{ij} p_{jl} + x_i {}_k\right) &= \left(\sum_{l=1}^{m} e_{lk} \sum_{j=1;j\neq i}^{n} w_{ij}\tilde{p}_{jl} + x_i \tilde{e}_k\right) \\ {}_k^2 &= e_k^2 \end{aligned} \tag{28}$$

And in matrix form, it will be

$$\begin{aligned} WP \quad + XB \ &= WP^{e} \ + X\tilde{B} \\ D_m \ &= \tilde{D}_m \end{aligned} \tag{29}$$

If $[WP; X; L]$ has full column rank, the observational equivalence of $\gamma^0; B^0; D_m{}^0$ and $\tilde{e}^0; \tilde{B}^0; \tilde{D}_m{}^0$ implies that

$$\gamma^0; B^0; D_m{}^0 = \tilde{e}^0; \tilde{B}^0; \tilde{D}_m{}^0 \tag{30}$$

means that the reduced form parameters can be identified. Therefore, the following assumption is essential to our econometric model identification.

Assumption 4. $[WP; X; L]$ has full column rank.

When the Assumption 4 holds, we will have the sufficient conditions for the identification of the reduced form parameters $\gamma^0; B^0; D_m{}^0$, where $\gamma = \alpha^{-1}$, $B = B\alpha^{-1}$, and D_m is an $m \times m$ diagonal matrix whose diagonal elements are α^{0-1} α^{-1} and zeros in non-diagonal positions. To figure out the sufficient condition for the identification of structural form parameters, we need to propose some constraints to the structural form matrix $\Gamma = [\gamma^0; \alpha^0; B^0]^0$. Suppose γ_k is the k-th column of Γ, and R_k is the matrix for the constraint that $R_k\gamma_k = 0$ and $\text{rank}(R_k) = m - 1$ for $k = 1; ; m - 1$, which is the sufficient rank condition to identify structural parameters from reduced form parameters by Schmidt (1976).

Assumption 5. Let $\Gamma = [\gamma^0; \alpha^0; B^0]^0$, and γ_k is the k-th column of Γ, and the R_k is the matrix for constraints $R_k\gamma_k = 0$, and $\text{rank}(R_k) = m - 1$ for $k = 1; ; m$.

Under all the assumptions above, we can derive a two-stage estimation process of the econometric model's structural parameters in the next section.

4 Estimation

We will derive the estimation of the model based on the identification process in the previous section. There are two steps of the estimation. The first step is to estimate the reduced form parameters γ, B, and $(\sigma_1^2; ; \sigma_m^2)$ by Nested Pseudo Likelihood (NPL) algorithm, which is discussed in Aguirregabiria and Mira (2007). NPL is also applied in Lin and Xu (2017) in large network games and adopted in Liu (2019) for multi-activity network games with discrete outcomes. Suppose we denote $\theta = [\gamma^0; B^0; L_m]^0$ and at $t = 0$ the NPL starts from an initial vector $p^{(0)} \in [0; +1)^{mn}$ and conduct the following iterative steps:

Step 1 Given $\mathbf{p}^{(t-1)}$, obtain $\hat{\theta}_k^{(t)} = (\hat{\lambda}_{1k}^{(t)}, \ldots, \hat{\lambda}_{mk}^{(t)}, \hat{\beta}_k^{(t)\prime}, \sigma_k^{(t)})' = \arg\max \ln L(\theta_k; \mathbf{p}^{(t-1)})$ where

$$\begin{aligned}\ln L(\theta_k; \mathbf{p}^{(t-1)}) &= \sum_{i=1}^{N} d_{ik} \ln \frac{1}{\sigma_k} \phi\left(\frac{y_{ik} - u_{ik}^{(t-1)}}{\sigma_k}\right) + (1 - d_{ik}) \left[1 - \Phi\left(\frac{u_{ik}^{(t-1)}}{\sigma_k}\right)\right] \\ &= \sum_{i=1}^{N} \left\{ d_{ik} \left[-\frac{1}{2}\ln 2\pi - \frac{1}{2}\ln \sigma_k^2 - \frac{1}{2\sigma_k^2}\left(y_{ik} - u_{ik}^{(t-1)}\right)^2 \right] + (1 - d_{ik}) \left[1 - \Phi\left(\frac{u_{ik}^{(t-1)}}{\sigma_k}\right)\right] \right\}\end{aligned}$$

where

$$u_{ik}^{(t-1)} = \sum_{l=1}^{m} \lambda_{lk} \sum_{j \neq i}^{n} w_{ij} p_{jl}^{(t-1)} + \mathbf{x}_i' \beta_k$$

for $k \in \{1, 2, \ldots, m\}$

Step 2 Given $\hat{\theta}^{(t)} = [\hat{\theta}_1^{(t)}, \ldots, \hat{\theta}_m^{(t)}]$, obtain $\mathbf{p}^{(t)} = \hat{h}(\mathbf{p}^{(t-1)}; \hat{\theta}^{(t)})$, where

$$\hat{h}(\mathbf{p}^{(t-1)}; \hat{\theta}^{(t)}) = [\hat{h}_1(\mathbf{p}^{(t-1)}; \hat{\theta}^{(t)})', \ldots, \hat{h}_m(\mathbf{p}^{(t-1)}; \hat{\theta}^{(t)})']'$$

with

$$\hat{h}_k(\mathbf{p}^{(t-1)}; \hat{\theta}^{(t)}) = \begin{array}{l} \sum_{l=1}^{m} \hat{\lambda}_{1k}^{(t)} \sum_{j=1}^{n} w_{1j} p_{jl}^{(t-1)} + x_1' \hat{\beta}_k^{(t)} \Phi\left(\dfrac{\sum_{l=1}^{m} \hat{\lambda}_{1k}^{(t)} \sum_{j=1}^{n} w_{1j} p_{jl}^{(t-1)} + x_1' \hat{\beta}_k^{(t)}}{\sigma_k^{(t)}}\right) \\ \quad + \sigma_k^{(t)} \phi\left(\dfrac{\sum_{l=1}^{m} \hat{\lambda}_{1k}^{(t)} \sum_{j=1}^{n} w_{1j} p_{jl}^{(t-1)} + x_1' \hat{\beta}_k^{(t)}}{\sigma_k^{(t)}}\right) \end{array}$$

Xu (2017), and Liu (2019). The detailed steps of deriving the asymptotic distribution of the NPL algorithm estimator are in Appendix.

Suppose we denote all the regressing variables as $Z = [WP; X]$, then we can rewrite our reduced form model as following

$$Y = Z[\ ^{0}; B\ ^{0}]^{0} \quad E = Z \qquad E \tag{31}$$

After the estimation of reduced form parameters , B , and based on the rank condition of constraints of the structural form parameters, we can estimate the reduced form parameters ,

then we can derive the reduced

(42)

the matrix form of red are

$$= \quad ^{-1} = \begin{bmatrix} 11 & 12 \\ 21 & 22 \end{bmatrix} \qquad B = B^{-1} = \qquad (43)$$

and the relation between reduced-form parameters and struc expressed as

$$_{21} = \frac{12}{11} \qquad _{12} = \frac{21}{22} \qquad _{1} = _{11} \quad \frac{12\ 21}{22}$$

$$_{11} = _{11} \quad \frac{21}{22}\ _{12} \qquad _{12} = _{12} \quad \frac{12}{11}\ _{11} \qquad _{22} = _{22} \quad \frac{12}{11}\ _{21} \qquad (44)$$

the equation (44) gives an arithmeti relation between reduced-form parameters and structural-form p we know the true value of the reduced-form parameters, we can use t calculate the true value of structural-form parameters. However, as w reduced-form parameters in our estimation and simulation, we need the f ur algorithm estimator to show the relation between the reduced-form esti d in the Appendix. As for the

(45)

$$_{1\ 2} \qquad _{2}$$

$$_{22} \quad \overset{n}{\geq}$$

what we want to estimate (; $_1$; $_2$) are correlation and standard deviation, which means that 2 (1; 1), $_1$ 2 (0; + 1), and $_2$ 2 (0; + 1), therefore, we introduce the following transformation during the optimization

$$_1 = f(a_1) = e^{a_1} \qquad _2 = f(a_2) = e^{a_2} \qquad = g(a_3) = 1 \quad \frac{2}{1 + e^{a_3}} \tag{47}$$

after this transformation, we can search a_1, a_2, and a_3 all along the whole real line. And for any function, the rst-order partial derivative with respect to a_1, a_2, and a_3 can be written as the rst-order partial derivative with respect to $_1$, $_2$, .

$$\begin{aligned}
\frac{@ \ln L}{@ a_1} &= \frac{@ \ln L}{@ _1}\frac{@ _1}{@ a_1} = {}_1\frac{@ \ln L}{@ _1} \\
\frac{@ \ln L}{@ a_2} &= \frac{@ \ln L}{@ _2}\frac{@ _2}{@ a_2} = {}_2\frac{@ \ln L}{@ _2} \\
\frac{@ \ln L}{@ a_3} &= \frac{@ \ln L}{@ }\frac{@ }{@ a_3} = \frac{@ \ln L}{@ }
\end{aligned} \tag{48}$$

The steps of the algorithm are the following.

At step 0, suppose we initialize $p_1^{(0)} = p_2^{(0)}$ 2 $[0; 1]^n$, and initialize $_{pq}^{(0)} = _{pq}^{(0)} = 0{:}1$,

At step $t \geq 1$ and we de ne $d_{i1} = 1(y_{i1} > 0)$, $d_{i2} = 1(y_{i2} > 0)$, then we can write the log-likelihood as

$$\begin{aligned}
\ln L_1 = {} & \sum_{i=1}^{} d_{i1} \ln \left[\frac{1}{_1} \quad \frac{_{11} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + _{21} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1} \ _{11} + x_{i2} \ _{21} \quad y_{i1}}{_1} \right] \\
& + (1 \quad d_{i1}) \ln \left[1 \quad \Phi \frac{_{11} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + _{21} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1} \ _{11} + x_{i2} \ _{21}}{_1} \right] \\
\ln L_2 = {} & \sum_{i=1}^{} d_{i2} \ln \left[\frac{1}{_2} \quad \frac{_{12} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + _{22} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1} \ _{12} + x_{i2} \ _{22} \quad y_{i2}}{_2} \right] \\
& + (1 \quad d_{i2}) \ln \left[1 \quad \Phi \frac{_{12} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + _{22} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1} \ _{12} + x_{i2} \ _{22}}{_2} \right]
\end{aligned} \tag{49}$$

the results are

$$\begin{aligned}
({}_{11}^{(t)}; {}_{21}^{(t)}; {}_{11}^{(t)}; {}_{21}^{(t)}; {}_{1}^{2(t)}) &= \arg\max \ln L_1(_{11}; _{21}; _{11}; _{21}; _1) \\
({}_{12}^{(t)}; {}_{22}^{(t)}; {}_{12}^{(t)}; {}_{22}^{(t)}; {}_{2}^{2(t)}) &= \arg\max \ln L_2(_{12}; _{22}; _{12}; _{22}; _2)
\end{aligned} \tag{50}$$

then we have the $\mathbf{p}_1^{(t+1)} = (p_{11}^{(t+1)}; \quad ; p_{n1})'$ and $\mathbf{p}_2^{(t+1)}(p_{12}^{(t+1)}; \quad ; p_{n2})'$ as

$$
\begin{aligned}
p_{i1}^{(t+1)} &= (\ {}_{11}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + \ {}_{21}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1}\ {}_{11}^{(t)} + x_{i2}\ {}_{21}^{(t)})\Phi_{i1}^{(t)} + \ {}_{1}^{(t)}\ {}_{i1}^{(t)} \\
p_{i2}^{(t+1)} &= (\ {}_{12}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + \ {}_{22}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1}\ {}_{12}^{(t)} + x_{i2}\ {}_{22}^{(t)})\Phi_{i2}^{(t)} + \ {}_{2}^{(t)}\ {}_{i2}^{(t)}
\end{aligned}
\tag{51}
$$

for $i = 1; \quad ; n$ and we have

$$
\begin{aligned}
\Phi_{i1}^{(t)} &= \Phi\ \frac{{}_{11}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + \ {}_{21}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1}\ {}_{11}^{(t)} + x_{i2}\ {}_{21}^{(t)}}{} \\
{}_{i1}^{(t)} &= \ \frac{{}_{11}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + \ {}_{21}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1}\ {}_{11}^{(t)} + x_{i2}\ {}_{21}^{(t)}}{{}^{1}} \\
\Phi_{i2}^{(t)} &= \Phi\ \frac{{}_{12}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + \ {}_{22}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1}\ {}_{12}^{(t)} + x_{i2}\ {}_{22}^{(t)}}{} \\
{}_{i2}^{(t)} &= \ \frac{{}_{12}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i1}^{(t)} + \ {}_{22}^{(t)} \sum_{j=1}^{n} w_{ij} p_{i2}^{(t)} + x_{i1}\ {}_{12}^{(t)} + x_{i2}\ {}_{22}^{(t)}}{{}^{2}}
\end{aligned}
\tag{52}
$$

the same as previous notation, $(\ )$ and $\Phi(\ )$ represent the PDF and CDF of the standard normal distribution. We repeat step t and $(t + 1)$ until the parameters estimated converge. Then, use the estimated parameters, respectively, $\hat{}_{11}; \hat{}_{12}; \hat{}_{21}; \hat{}_{22}; \hat{}_{11}; \hat{}_{12}; \hat{}_{21}; \hat{}_{22}$ and $\hat{}_1; \hat{}_2$, calculated the equilibrium, $\hat{p}_1$ and $\hat{p}_2$; then use all these to estimate $\quad$, the log-likelihood function is

$$
\begin{aligned}
\ln L(\ ) = \sum_{i=1}^{n} \Big\{ & d_{i1} d_{i2} \ln\ {}_{2}\ \frac{\hat{}_{11} \sum_{j=1}^{n} w_{ij} \hat{p}_{i1} + \hat{}_{21} \sum_{j=1}^{n} w_{ij} \hat{p}_{i2} + x_{i1} \hat{}_{11} + x_{i2} \hat{}_{21} \quad y_{i1}}{\hat{}_1}; \\
& \frac{\hat{}_{12} \sum_{j=1}^{n} w_{ij} \hat{p}_{i1} + \hat{}_{22} \sum_{j=1}^{n} w_{ij} \hat{p}_{i2} + x_{i1} \hat{}_{12} + x_{i2} \hat{}_{22} \quad y_{i2}}{\hat{}_2}; \hat{}_1; \hat{}_2; \\
& + d_{i1}(1 \quad d_{i2}) \ln \Phi\ \frac{(\hat{}_{11} \sum_{j=1}^{n} w_{ij} \hat{p}_{i1} + \hat{}_{21} \sum_{j=1}^{n} w_{ij} \hat{p}_{i2} + x_{i1} \hat{}_{11} + x_{i2} \hat{}_{21} \quad y_{i1})}{\hat{}_1 \sqrt{1 \quad {}_{2}}} \\
& \frac{\hat{}_{12} \sum_{j=1}^{n} w_{ij} \hat{p}_{i1} + \hat{}_{22} \sum_{j=1}^{n} w_{ij} \hat{p}_{i2} + x_{i1} \hat{}_{12} + x_{i2} \hat{}_{22}}{\hat{}_2 \sqrt{1 \quad {}_{2}}} \\
& \frac{\hat{}_{11} \sum_{j=1}^{n} w_{ij} \hat{p}_{i1} + \hat{}_{21} \sum_{j=1}^{n} w_{ij} \hat{p}_{i2} + x_{i1} \hat{}_{11} + x_{i2} \hat{}_{21} \quad y_{i1}}{\hat{}_1} \Big] \\
& + (1 \quad d_{i1}) d_{i2} \ln \Phi\ \frac{(\hat{}_{12} \sum_{j=1}^{n} w_{ij} \hat{p}_{i1} + \hat{}_{22} \sum_{j=1}^{n} w_{ij} \hat{p}_{i2} + x_{i1} \hat{}_{12} + x_{i2} \hat{}_{22} \quad y_{i2})}{\hat{}_2 \sqrt{1 \quad {}_{2}}}
\end{aligned}
$$

$$\frac{\hat{\alpha}_{11}\sum_{j=1}^{n} w_{ij}\hat{p}_{i1} + \hat{\alpha}_{21}\sum_{j=1}^{n} w_{ij}\hat{p}_{i2} + x_{i1}\hat{\alpha}_{11} + x_{i2}\hat{\alpha}_{21}}{\hat{\sigma}_1\sqrt{1-\hat{\rho}^2}}$$

$$\left.\frac{\hat{\alpha}_{12}\sum_{j=1}^{n} w_{ij}\hat{p}_{i1} + \hat{\alpha}_{22}\sum_{j=1}^{n} w_{ij}\hat{p}_{i2} + x_{i1}\hat{\alpha}_{12} + x_{i2}\hat{\alpha}_{22} - y_{i2}}{\hat{\sigma}_2}\right]$$

$$+(1-d_{i1})(1-d_{i2})\ln\Phi_2\left[\frac{\hat{\alpha}_{11}\sum_{j=1}^{n} w_{ij}\hat{p}_{i1} + \hat{\alpha}_{21}\sum_{j=1}^{n} w_{ij}\hat{p}_{i2} + x_{i1}\hat{\alpha}_{11} + x_{i2}\hat{\alpha}_{21}}{\hat{\sigma}_1}, \right.$$

$$\left.\left.\frac{\hat{\alpha}_{12}\sum_{j=1}^{n} w_{ij}\hat{p}_{i1} + \hat{\alpha}_{22}\sum_{j=1}^{n} w_{ij}\hat{p}_{i2} + x_{i1}\hat{\alpha}_{12} + x_{i2}\hat{\alpha}_{22}}{\hat{\sigma}_2}; \hat{\sigma}_1, \hat{\sigma}_2\right]\right]$$

where $_2(\ ;\ ;\hat{\sigma}_1;\hat{\sigma}_2;\ )$ and $\Phi_2(\ ;\ ;\hat{\sigma}_1;\hat{\sigma}_2;\ )$ are PDF and CDF of bivariate normal distributed random variables with variance-covariance matrix

$$\begin{bmatrix} \hat{\sigma}_1^2 & \hat{\sigma}_1\hat{\sigma}_2 \\ \hat{\sigma}_1\hat{\sigma}_2 & \hat{\sigma}_2^2 \end{bmatrix} \tag{53}$$

and the estimation result of , i.e., $\hat{}$ can be written as

$$\hat{} = \arg\max \ln L(\ ) \tag{54}$$

Suppose we denote $Z = [W\mathbf{p_1}, W\mathbf{p_2}, X]$, and $_1 = (\ _{11};\ _{21};\ _{11};\ _{21})'$ and $_2 = (\ _{12};\ _{22};\ _{12};\ _{22})'$, then we have

$$\begin{aligned} y_1 &= Z\ _1\ \ _1 \\ y_2 &= Z\ _2\ \ _2 \end{aligned} \tag{55}$$

then given the NPL algorithm, for each step, suppose the current equilibrium $\hat{p} = (\hat{p}_1', \hat{p}_2')'$, and we use $\hat{Z} = [W\hat{p}_1, W\hat{p}_2, X]$, then we have

$$\begin{aligned} (\hat{}_1;\hat{}_1^2) &= \arg\max \ln L(\ _1;\ _1^2;\hat{p}_1) \\ (\hat{}_2;\hat{}_2^2) &= \arg\max \ln L(\ _2;\ _2^2;\hat{p}_2) \end{aligned} \tag{56}$$

where

$$\begin{aligned} \ln L(\ _1;\ _1;\hat{p}_1) &= \sum_{i=1} d_{i1}\ln\left[\frac{1}{\ _1}\ \ \frac{\hat{z}_i'\ _1 - y_{i1}}{\ _1}\right] + (1-d_{i1})\ln\left[1-\Phi\left[\frac{\hat{z}_i'\ _1}{\ _1}\right]\right] \\ \ln L(\ _2;\ _2;\hat{p}_2) &= \sum_{i=1} d_{i2}\ln\left[\frac{1}{\ _2}\ \ \frac{\hat{z}_i'\ _2 - y_{i2}}{\ _2}\right] + (1-d_{i2})\ln\left[1-\Phi\left[\frac{\hat{z}_i'\ _2}{\ _2}\right]\right] \end{aligned} \tag{57}$$

the first-order conditions are

$$\begin{aligned}\frac{\partial L}{\partial \beta_k} &= \sum_{i=1}^{N} d_{ik}\frac{(z_i'\beta_k - y_{ik})z_i}{\sigma_k^2} - (1-d_{ik})\frac{\lambda_{ik}z_i}{(1-\Phi_{ik})(\sigma_k^2)^{\frac{1}{2}}} = 0 \\ \frac{\partial L}{\partial(\sigma_k^2)} &= \sum_{i=1}^{N} d_{ik}\left[-\frac{1}{2\sigma_k^2} + \frac{(z_i'\beta_k - y_{ik})^2}{2(\sigma_k^2)^2}\right] + (1-d_{ik})\frac{(z_i'\beta_k)\lambda_{ik}}{2(1-\Phi_{ik})(\sigma_k^2)^{\frac{3}{2}}} = 0\end{aligned} \tag{58}$$

where $k = \{1,2\}$, $\Phi_{ik} = \Phi(z_i'\beta_k/\sigma_k)$ and $\lambda_{ik} = \lambda(z_i'\beta_k/\sigma_k)$ then according to the results $E(y_{ik} - p_{ik}) = 0$, i.e.,

$$\begin{aligned}E[y_{ik} - (z_i'\beta_k)\Phi_{ik} - \sigma_k\lambda_{ik}] &= 0 \\ E[d_{ik} - \Phi_{ik}] &= 0\end{aligned} \tag{59}$$

this can be used to simplify the first-Taylor expansion of the first-order condition, which is the approach to derive the asymptotic variance-covariance matrix of reduce-form parameters' estimator discussed in Amemiya (1973), Amemiya (1985), and Maddala (1986). Detailed steps for deriving the variance-covariance matrix will be in the Appendix.

Remark 1. (Binary Dependent Variable Case) Liu (2019) proposes the case in which all the decision outcomes are binary. There is no need to estimate the variance of reduced form error terms σ_1 and σ_2. Therefore, the first-order conditions of the NPL estimator degenerate to

$$\frac{\partial \ln L(\hat{\beta}_k, \hat{p})}{\partial \beta_k} = \sum_{i=1}^{N}\frac{[d_{ik} - \Phi(\hat{z}_i'\beta_k)]}{\Phi(\hat{z}_i'\beta_k)[1-\Phi(\hat{z}_i'\beta_k)]}\hat{z}_i = 0 \tag{60}$$

for $k = 1,2$. The first-order Taylor expansion of the above equation around the can draw the following equation

$$\begin{aligned}&\sum_{i=1}^{N}\frac{(d_{ik}-\Phi_{ik})\lambda_{ik}}{\Phi_{ik}(1-\Phi_{ik})}z_i - \frac{(d_{ik}-\Phi_{ik})\lambda_{ik}}{\Phi_{ik}(1-\Phi_{ik})}z_i(z_i'\beta_k)z_i' + \sigma_{1k}w_i\frac{\partial p_1}{\partial \beta_k'} + \sigma_{2k}w_i\frac{\partial p_2}{\partial \beta_k'}(\hat{\beta}_k - \beta_k) \\ &\quad - \frac{\lambda_{ik}^2}{\Phi_{ik}(1-\Phi_{ik})}z_i z_i' + \sigma_{1k}w_i\frac{\partial p_1}{\partial \beta_k'} + \sigma_{2k}w_i\frac{\partial p_2}{\partial \beta_k'}(\hat{\beta}_k - \beta_k) \\ &\quad + \frac{(d_{ik}-\Phi_{ik})\lambda_{ik}}{\Phi_{ik}(1-\Phi_{ik})}0' - w_i\frac{\partial p_1}{\partial \beta_k'} - w_i\frac{\partial p_2}{\partial \beta_k'}(\hat{\beta}_k - \beta_k) \\ &\quad - \frac{(d_{ik}-\Phi_{ik})\lambda_{ik}^2(1-2\Phi_{ik})}{\Phi_{ik}(1-\Phi_{ik})}z_i z_i' + \sigma_{1k}w_i\frac{\partial p_1}{\partial \beta_k'} + \sigma_{2k}w_i\frac{\partial p_2}{\partial \beta_k'}(\hat{\beta}_k - \beta_k) \\ =\ &\sum_{i=1}^{N}\frac{(d_{ik}-\Phi_{ik})\lambda_{ik}}{\Phi_{ik}(1-\Phi_{ik})}z_i - \frac{\lambda_{ik}^2}{\Phi_{ik}(1-\Phi_{ik})}z_i z_i' + \sigma_{1k}w_i\frac{\partial p_1}{\partial \beta_k'} + \sigma_{2k}w_i\frac{\partial p_2}{\partial \beta_k'}(\hat{\beta}_k - \beta_k)\end{aligned}$$

$= 0zzzzz$

$= O_p(1)$

where $\Phi_{ik} = \Phi(z_i^0 \ _k)$ and $\ _{ik} = \ (z_i^0 \ _k)$

5.2 Simulation Results

We conduct two types of normalized network structures in our simulation. The rst type is the random network, which means agent i will be randomly a ected by ve other agents in the network. And each of the ve agents' e ects is identical. In this case, that is $w_{ij} = 1=5$ if agent j can a ect agent i in the network. The other type is the circular network structure. In this case, the agent i will only connect with agent $i+1$ and i 1, and the peer e ects are the same, i.e., $w_{i;i+1} = w_{i;i-1} = 1=2$. And for agent 1, we have $w_{12} = w_{1n} = 1=2$. And for agent n, we have $w_{n1} = w_{n;n-1} = 1=2$. The network graph is similar to a big circle in which each node only connects its two neighbors. That is where the network name `circular' comes from. The parameter of the simulation are followings, $_{12} = \ _{21} = 0:5$, $\ _1 = \ _2 = 1$, $n = 2000$, and rep = 1000, $\ _1 = \ _2 = 1$, $\ _{12} = 0:1$.

All the detailed simulation results are available in part B of the Appendix. In the following discussion, we will focus on three typical scenarios

Case 1: Weak peer e ect $\ _{11} = \ _{22} = 0:2$, $\ _{12} = \ _{21} = 0:1$

? Random Network

Case 2: Medium peer e ect $_{11} = {}_{22} = 0{:}5$, $_{12} = {}_{21} = 0{:}3$

? Random Network

$\hat{}_{12}$	$\hat{}_{21}$	$\hat{}_{11}$	$\hat{}_{12}$	$\hat{}_{21}$	$\hat{}_{22}$
0.500	0.502	0.542	0.329	0.332	0.546
(0.031)	(0.031)	(0.053)	(0.056)	(0.059)	(0.053)
$\hat{}_1$	$\hat{}_2$	$\hat{}_1$	$\hat{}_2$	$\hat{}_{12}$	
1.000	0.999	0.998	0.996	0.099	
(0.047)	(0.047)	(0.029)	(0.029)	(0.029)	

(0.029)

? Cie4)

0.546

From the results, we could nd that the random network estimation of simultaneous e ect matrix parameters ($_{12}$ and $_{11}$) performs better than those in circular networks. In a random network, the estimator average accuracy of simultaneous e ect matrix parameters ($_{12}$ and $_{11}$) is always low, and the coe cient of variation is about 6% 7%. However, in a circular network, the estimator average accuracy of the simultaneous e ect matrix parameter can be lower than 95%, and the coe cient of variation can be over 8%.

As the peer e ect strengthens, the estimator's accuracy and consistency increase. In a strong peer e ect case, the estimation average accuracy of the same-activity peer e ect ($_{11}$ and $_{22}$) is over 98%, and the coe cient of variation is about 2% in a circular network and 4% in a random network. As for the cross-activity peer e ect, the accuracy is over 95%, and the coe cient of variation is about 10%. Both perspectives show under a strong peer e ect case, the estimator works better.

6 Conclusion

This paper derives a simultaneous equation model with peer e ects and rational expectations under an incomplete information network. The econometric model and the microeconomic foundation are discussed. The su cient condition of the existence of a Bayesian Nash Equilibrium under an incomplete information network game is derived. A nested pseudo-likelihood estimation process and the estimator's asymptotic distribution properties are developed. Monte Carlo simulation shows the consistency of the estimation results when the sample size is nite and the network size is large. In this current paper, the network is predetermined and has no relation with agents' random shock and attributes. However, in a real economic situation, an agent's connection with other agents in the network can partially re ect his/her observed and unobserved characteristics. Therefore, the endogenous network structure can be a potential future extension. Another extension in our current work is that our econometric model proposes the simultaneous e ect from the intention of di erent activities of each agent. However, it is very possible that the simultaneous e ect is the relation between the activity's intention and other activities' outcomes. This means the condition of an option model is necessa e(is)-330(necr)] istts

References

Aguirregabiria, V. and P. Mira (2007). Sequential estimation of dynamic discrete games. Econometrica 75(1), 1{53.

Amemiya, T. (1973). Regression analysis when the dependent variable is truncated normal. Econometrica: Journal of the Econometric Society, 997{1016.

Amemiya, T. (1974). Multivariate regression and simultaneous equation models when the dependent variables are truncated normal. Econometrica: Journal of the Econometric Society, 999{1012.

Amemiya, T. (1978). The estimation of a simultaneous equation generalized probit model. Econometrica: Journal of the Econometric Society, 1193{1205.

Amemiya, T. (1979). The estimation of a simultaneous-equation tobit model. International economic review, 169{181.

Amemiya, T. (1985). Advanced Econometrics. Harvard university press.

Ballester, C., A. Calvo-Armengol, and Y. Zenou (2006). Who's who in networks. wanted: The key player. Econometrica 74(5), 1403{1417.

Blume, L. E., W. A. Brock, S. N. Durlauf, and R. Jayaraman (2015). Linear social interactions models. Journal of Political Economy 123(2), 444{496.

Calvo-Armengol, A., E. Patacchini, and Y. Zenou (2009). Peer e ects and social networks in education. The review of economic studies 76(4), 1239{1267.

Cohen-Cole, E., X. Liu, and Y. Zenou (2018). Multivariate choices and identi cation of social interactions. Journal of Applied Econometrics 33(2), 165{178.

Horn, R. A. and C. R. Johnson (2012). Matrix analysis. Cambridge university press.

Kasahara, H. and K. Shimotsu (2012). Sequential estimation of structural models with a xed point constraint. Econometrica 80(5), 2303{2319.

Xu, X. and L.-f. Lee (2015). Maximum likelihood estimation of a spatial autoregressive tobit model. Journal of Econometrics 188 (1), 264{280.

Yang, C. and L.-f. Lee (2017). Social interactions under incomplete information with heterogeneous expectations. Journal of Econometrics 198 (1), 65{83.

Yang, C., L.-f. Lee, and X. Qu (2018). Tobit models with social interactions: Complete vs incomplete information. Regional Science and Urban Economics 73, 30{50.

Appendices

A. AGLS Estimator

Let $m = 2$, according to Lee (1978, 1979, 1981, 1982), Maddala and Lee (1976), Nelson and Olson (1978), and Liu (2019), we have the structural model as

$$\begin{aligned} y_1 &= y_2 \,{}_{21} + Z_1 \,{}_1 \quad {}_1 \\ y_2 &= y_1 \,{}_{12} + Z_2 \,{}_2 \quad {}_2 \end{aligned} \tag{61}$$

where $Z_1 = [Wp_1; Wp_2; X_1]$, $Z_2 = [Wp_1; Wp_2; X_2]$, and $Z = [Wp_1; Wp_2; X]$. Then, we can write the reduced form as

$$\begin{aligned} y_1 &= Z \,{}_1 \quad {}_1 \\ y_2 &= Z \,{}_2 \quad {}_2 \end{aligned} \tag{62}$$

For agent i, $z_i^{\prime} = (w_i p_1; w_i p_2; x_i^{\prime})$ is the i-th row of Z

Now, suppose we introduce the vectors of ones and zeros to form the selection matrices J_1 and J_2, subject to $Z_1 = ZJ_1$ and $Z_2 = ZJ_2$, we can rewrite the structural model as

$$\begin{aligned} y_1 &= y_2 \ _{21} + ZJ_1 \ _1 \quad _1 \\ y_2 &= y_1 \ _{12} + ZJ_2 \ _2 \quad _2 \end{aligned} \tag{65}$$

then we put the reduced-form equations in the model (62) into the reorganized structural model (65), we can get

$$\begin{aligned} y_1 &= (Z \ _1 \quad _1) \ _{21} + ZJ_1 \ _1 \quad _1 = Z(\quad _1 \ _{21} + J_1 \ _1) + \ _1 \ _{21} \quad _1 \\ y_2 &= (Z \ _2 \quad _2) \ _{12} + ZJ_2 \ _2 \quad _2 = Z(\quad _2 \ _{12} + J_2 \ _2) + \ _2 \ _{12} \quad _2 \end{aligned} \tag{66}$$

then, we can derive the relation between reduced-form parameters and structural form parameters as following

$$\begin{aligned} _1 &= \quad _2 \ _{21} + J_1 \ _1 \\ _2 &= \quad _1 \ _{12} + J_2 \ _2 \end{aligned} \tag{67}$$

then, we can derive the relation between the estimation of reduced-form parameters and the true value of structural parameters as following

$$\begin{aligned} b_1 &= b_2 \ _{21} + J_1 \ _1 + v_1 \\ b_2 &= b_1 \ _{12} + J_2 \ _2 + v_2 \end{aligned} \tag{68}$$

where

$$\begin{aligned} v_1 &= (b_1 \quad _1) + (b_2 \quad _2) \ _{21} \\ v_2 &= (b_2 \quad _2) + (b_1 \quad _1) \ _{12} \end{aligned} \tag{69}$$

Suppose we apply $\quad$ as the notation of the asymptotic covariance matrix of $v = (v_1^0; v_2^0)^0$. Suppose we have $_1 = (\ _{21};\ _1^0)^0$ and $_2 = (\ _{12};\ _2^0)^0$. Then the estimator $\quad = (\ _1^0;\ _2^0)^0$ is

$$b \ = \ (\Pi^0 b^{-1} \Pi)^{-1} \Pi^0 b^{-1} b \tag{70}$$

where

$$\hat{H} = \begin{bmatrix} \hat{H}_1 & 0 \\ 0 & \hat{H}_2 \end{bmatrix} \tag{71}$$

in which $\hat{H}_1 = [\ \ b_2; J_1]$ and $\hat{H}_2 = [\ \ b_1; J_2]$. And b is a consistent estimator of . To derive the detailed form of , we need to derive the asymptotic variance-covariance3(the)e7927(e)-313(need)-03Hce-c2

$$\frac{@}{@_k^2}\ \frac{@L}{@_k} = \sum_i$$

$$
\begin{aligned}
\frac{@}{@\ _k}\ \frac{@L}{@\ _k^2} &= \sum_{i=1}^{N} d_{ik}\frac{(z_i^0{}_k - y_{ik})}{(\ _k^2)^2} + (1 - d_{ik})\frac{{}_{ik}}{2(1 - \Phi_{ik})(\ _k^2)^{\frac{3}{2}}} \\
&\quad (1 - d_{ik})\frac{(z_i^0{}_k)^2\ {}_{ik}}{2(1 - \Phi_{ik})(\ _k^2)^{\frac{5}{2}}} + (1 - d_{ik})\frac{(z_i^0{}_k)\ {}^2_{ik}}{2(1 - \Phi_{ik})^2(\ _k^2)^2} \\
&\quad z_i + {}_{1k}\frac{@p_1^0}{@\ _k}w_i^0 + {}_{2k}\frac{@p_2^0}{@\ _k}w_i^0 \\
&= \sum_{i=1}^{N} d_{ik}\frac{(z_i^0{}_k - y_{ik})z_i}{(\ _k^2)^2} + (1 - d_{ik})\frac{z_i\ {}_{ik}}{2(1 - \Phi_{ik})(\ _k^2)^{\frac{3}{2}}} \\
&\quad (1 - d_{ik})\frac{(z_i^0{}_k)^2\ {}_{ik}z_i}{2(1 - \Phi_{ik})(\ _k^2)^{\frac{5}{2}}} + (1 - d_{ik})\frac{(z_i^0{}_k)\ {}^2_{ik}z_i}{2(1 - \Phi_{ik})^2(\ _k^2)^2} \\
&\quad + d_{ik}\frac{(z_i^0{}_k - y_{ik})}{(\ _k^2)^2} + (1 - d_{ik})\frac{{}_{ik}}{2(1 - \Phi_{ik})(\ _k^2)^{\frac{3}{2}}} \\
&\quad (1 - d_{ik})\frac{(z_i^0{}_k)^2\ {}_{ik}}{2(1 - \Phi_{ik})(\ _k^2)^{\frac{5}{2}}} + (1 - d_{ik})\frac{(z_i^0{}_k)\ {}^2_{ik}}{2(1 - \Phi_{ik})^2(\ _k^2)^2} \\
&\quad {}_{1k}\frac{@p_1^0}{@\ _k}w_i^0 + {}_{2k}\frac{@p_2^0}{@\ _k}w_i^0 \\
&= \sum_{i=1}^{N} \frac{z_i\ {}_{ik}}{2(\ _k^2)^{\frac{3}{2}}} - \frac{(z_i^0{}_k)^2\ {}_{ik}z_i}{2(\ _k^2)^{\frac{5}{2}}} + \frac{(z_i^0{}_k)\ {}^2_{ik}z_i}{2(1 - \Phi_{ik})(\ _k^2)^2} \\
&\quad + \frac{\Phi_{ik}(z_i^0{}_k - y_{ik})}{(\ _k^2)^2} + \frac{{}_{ik}}{2(\ _k^2)^{\frac{3}{2}}} - \frac{(z_i^0{}_k)^2\ {}_{ik}}{2(\ _k^2)^{\frac{5}{2}}} + \frac{(z_i^0{}_k)\ {}^2_{ik}}{2(1 - \Phi_{ik})(\ _k^2)^2} \\
&\quad {}_{1k}\frac{@p_1^0}{@\ _k}w_i^0 + {}_{2k}\frac{@p_2^0}{@\ _k}w_i^0
\end{aligned}
\tag{78}
$$

According to $p_{ik} = (x_i^{\prime}\ _k)\Phi_{ik} +\ _k\ _{ik}$, we can derive $@p_k=@\ _l$ and $@p_k=@\ _l{}^2$. ($k = 1;2$ and $l = 1;2$)

$$
\begin{aligned}
\frac{@p_1}{@\ _1^{\prime}} &= K_1^{-1}A_1Z \\
\frac{@p_2}{@\ _2^{\prime}} &= K_2^{-1}A_2Z \\
\frac{@p_1}{@\ _2^{\prime}} &= (I_n \quad {}_{11}A_1W)\ {}_{21}A_1WK_2^{-1}A_2Z \\
\frac{@p_2}{@\ _1^{\prime}} &= (I_n \quad {}_{22}A_2W)\ {}_{12}A_2WK_1^{-1}A_1Z \\
\frac{@p_1}{@\ _1{}^2} &= K_1^{-1}\frac{1}{2(\ _1{}^2)^{1=2}} \\
\frac{@p_2}{@\ _2{}^2} &= K_2^{-1}\frac{2}{2(\ _2{}^2)^{1=2}} \\
\frac{@p_1}{@\ _2{}^2} &= (I_n \quad {}_{11}A_1W)\ {}_{21}A_1WK_2^{-1}\frac{2}{2(\ _2{}^2)^{1=2}} \\
\frac{@p_2}{@\ _1{}^2} &= (I_n \quad {}_{22}A_2W)\ {}_{12}A_2WK_1^{-1}\frac{1}{2(\ _1{}^2)^{1=2}}
\end{aligned}
\tag{79}
$$

where

$$
\begin{aligned}
K_1 &= I_n \quad {}_{11}A_1W \quad {}_{12}\ {}_{21}A_1W(I_n \quad {}_{22}A_2W)^{-1}A_2W \\
K_2 &= I_n \quad {}_{22}A_2W \quad {}_{12}\ {}_{21}A_2W(I_n \quad {}_{11}A_1W)^{-1}A_1W \\
A_1 &= \mathrm{diag}(\Phi_{11};\Phi_{21};\quad \Phi_{n1}) \\
A_2 &= \mathrm{diag}(\Phi_{12};\Phi_{22};\quad \Phi_{n2})
\end{aligned}
\tag{80}
$$

Then we can derive $@(@\ln L=@\ _k)=@\ _k$, $@(@\ln L=@\ _k)=@\ _k{}^2$, $@(@\ln L=@\ _k{}^2)=@\ _k$, and $@(@\ln L=@\ _k{}^2)=@\ _k{}^2$ by previous results. According to the algebra results, we can derive the asymptotic variance of $(b_1^{\prime};b_1{}^2;b_2^{\prime};b_2{}^2)^{\prime}$ is

$$
V = \begin{bmatrix} V_{11} & V_{12} \\ V_{12}^{\prime} & V_{22} \end{bmatrix}
\tag{81}
$$

as for the diagonal element of V, i.e., V_{kk}, can be derived as following

$$
V_{kk} = \begin{bmatrix} A_{kk} & b_{kk} \\ c_{kk}^{\prime} & g_{kk} \end{bmatrix}^{-1} \begin{bmatrix} H_{kk} & p_{kk} \\ q_{kk}^{\prime} & s_{kk} \end{bmatrix} \begin{bmatrix} A_{kk} & b_{kk} \\ c_{kk}^{\prime} & g_{kk} \end{bmatrix}^{\prime-1}
\tag{82}
$$

all the elements to derive V

and the elements H_{kl}, p_{kl}, q_{kl}, and s_{kl} can be derived as following

$$
\begin{aligned}
H_{kl} = Z^{0} \ \mathrm{diag}_{i=1}^{n} \ & \frac{(z_i^0{}_k - y_{ik})(z_i^0{}_l - y_{il})}{{}_k^2 \, {}_l^2}\Phi_2 \ \frac{z_i^0{}_k}{k}; \frac{z_i^0{}_l}{l}; \\
& + \frac{(z_i^0{}_k - y_{ik})}{(1 - \Phi_{il}) \, {}_k^2 \, {}_l}\Phi_2 \ \frac{z_i^0{}_k}{k}; \ \frac{z_i^0{}_l}{l}; \\
& + \frac{(z_i^0{}_l - y_{il})}{(1 - \Phi_{ik}) \, {}_k \, {}_l^2}\Phi_2 \ \frac{z_i^0{}_k}{k}; \frac{z_i^0{}_l}{l}; \\
& + \frac{}{(1 - \Phi_{ik})(1 - \Phi_{il}) \, {}_k \, {}_l} \ \Phi_2 \ \frac{z_i^0{}_k}{k}; \ \frac{z_i^0{}_l}{l}; \ Z
\end{aligned}
\tag{91}
$$

$$
\begin{aligned}
p_{kl} = \ & Z^{0}\mathrm{diag}_{i=1}^{n} \ \frac{(z_i^0{}_k - y_{ik})}{{}_k^2} \ \frac{(z_i^0{}_l - y_{il})^2}{2({}_l^2)^2} \ \frac{1}{2 \, {}_l^2} \ \Phi_2 \ \frac{z_i^0{}_k}{k}; \frac{z_i^0{}_l}{l}; \\
& + \frac{(z_i^0{}_k - y_{ik})(z_i^0{}_l)}{2(1 - \Phi_{il}) \, {}_k^2({}_l^2)^{\frac{3}{2}}}\Phi_2 \ \frac{z_i^0{}_k}{k}; \ \frac{z_i^0{}_l}{l}; \\
& + \frac{}{(1 - \Phi_{ik}) \, {}_k} \ \frac{(z_i^0{}_l - y_{il})^2}{2({}_l^2)^2} \ \frac{1}{2 \, {}_l^2} \ \Phi_2 \ \frac{z_i^0{}_k}{k}; \frac{z_i^0{}_l}{l}; \\
& + \frac{(z_i^0{}_l)}{2(1 - \Phi_{ik})(1 - \Phi_{il}) \, {}_k({}_l^2)^{\frac{3}{2}}}\Phi_2 \ \frac{z_i^0{}_k}{k}; \ \frac{z_i^0{}_l}{l}; \ {}_n
\end{aligned}
\tag{92}
$$

$$
\begin{aligned}
q_{kl} = \ & Z^{0}\mathrm{diag}_{i=1}^{n} \ \frac{(z_i^0{}_l - y_{il})}{{}_l^2} \ \frac{(z_i^0{}_k - y_{ik})^2}{2({}_k^2)^2} \ \frac{1}{2 \, {}_k^2} \ \Phi_2 \ \frac{z_i^0{}_k}{k}; \frac{z_i^0{}_l}{l}; \\
& + \frac{}{(1 - \Phi_{il}) \, {}_l} \ \frac{(z_i^0{}_k - y_{ik})^2}{2({}_k^2)^2} \ \frac{1}{2 \, {}_k^2} \ \Phi_2 \ \frac{z_i^0{}_k}{k}; \ \frac{z_i^0{}_l}{l}; \\
& + \frac{(z_i^0{}_l - y_{il})(z_i^0{}_k)}{2(1 - \Phi_{ik}) \, {}_l^2({}_k^2)^{\frac{3}{2}}}\Phi_2 \ \frac{z_i^0{}_k}{k}; \frac{z_i^0{}_l}{l}; \\
& + \frac{(z_i^0{}_k)}{2(1 - \Phi_{ik})(1 - \Phi_{il}) \, {}_l({}_k^2)^{\frac{3}{2}}}\Phi_2 \ \frac{z_i^0{}_k}{k}; \ \frac{z_i^0{}_l}{l}; \ {}_{2\,n}
\end{aligned}
\tag{93}
$$

$$
s_{kl} = {}_n^0\mathrm{diag}_{i=1}^{n} \ \frac{(z_i^0{}_k - y_{ik})^2}{2({}_k^2)^2} \ \frac{1}{2 \, {}_k^2} \ \frac{(z_i^0{}_l - y_{il})^2}{2({}_l^2)^2} \ 1_l{}^2_k) \ \Phi_2
$$