

## Geometry of epitaxial GaAs/(Al,Ga)As quantum dots as seen by excitonic spectroscopy

Jun-Wei Luo\*

*Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300, Republic of China*

80401, .

Alex Zunger†

*Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300, Republic of China*

80309, .

(Received 11 October 2011; published 28 December 2011)

It is shown that exciton and multiexciton emission lines (“spectral barcode”) of a quantum dot conceal nontrivial structural information on the shape and size of the dot, information which can be uncovered by comparison with





leading to the misuse of the FSS to infer shape anisotropy: In the Luttinger Hamiltonian representation, the effective mass of hole is anisotropic in that its value along (100) is different from along (110). Thus, if one ignores the fact that the QDs under consideration are made of atomistically discrete materials, the symmetry of circular based dot in this Hamiltonian is  $C_{4v}$ . Despite this, numerous papers<sup>18,31</sup> claimed that circular-based lens shape dot has  $D_{2d}$  symmetry. This is because in a continuum approximation the [110] and  $[1\bar{1}0]$  directions are equivalent. In such a  $D_{2d}$  symmetry, the fourfold degenerate exciton (originating from an electron of  $J_z = \pm 1/2$  and a heavy-hole of  $J_z = \pm 3/2$ ) splits into double-degenerate bright state ( $\Gamma_5$ ) and two nondegenerate dark states ( $\Gamma_1$  and  $\Gamma_2$ , respectively). Because  $\Gamma_5$  is degenerate in this approximation, the FSS is zero for cylindrically symmetric dots under the continuum point of view.

To account for the observed nonzero FSS, the continuum theory assumes that the FSS originates, in its entirety, from deviations from cylindrical symmetry of the overall QD shape, i.e., shape anisotropy of the QDs.<sup>17,19,20,30</sup> This shape anisotropy (e.g., elongation in  $[1\bar{1}0]$  direction<sup>17,19,20</sup>) of QD lowers then the  $D_{2d}$  symmetry to  $C_{2v}$ .<sup>32</sup> The double-degenerate bright  $\Gamma_5$  further splits into two nondegenerate states ( $\Gamma_2$  and  $\Gamma_4$ ). The lifting of the degeneracy of the two bright exciton states is referred to FSS and is used under the continuum point of view to fit the measured FSS into a geometric shape anisotropy.

In reality, the [110] and  $[1\bar{1}0]$  directions are nonequivalent in zincblende crystal. This leads to  $91([1])TJti$

## ACKNOWLEDGMENTS

This work was funded by the US Department of Energy, Office of Science, Basic Energy Science, Materials Sciences

and Engineering, under Contract No. DE-AC36-08GO28308 to NREL.

\*jun-wei.luo@nrel.gov

†alex.zunger@gmail.com

<sup>1</sup>G. M. Barrow, *J. Appl. Phys.* **52**, 1047 (1982). (McGraw-Hill, New York, 1962).

<sup>2</sup>M. Abbarchi, T. Kuroda, T. Mano, K. Sakoda, C. A. Mastrandrea, A. Vinattieri, M. Gurioli, and T. Tsuchiya, *Phys. Rev. B* **82**, 201301(R) (2010).

<sup>3</sup>M. Ediger, G. Bester, A. Badolato, P. M. Petroff, K. Karrai, A. Zunger, and R. J. Warburton, *Nat. Phys.* **3**, 774 (2007).

<sup>4</sup>E. Poem, J. Shemesh, I. Marderfeld, D. Galushko, N. Akopian, D. Gershoni, B. D. Gerardot, A. Badolato, and P. M. Petroff, *Phys. Rev. B* **76**, 235304 (2007).

<sup>5</sup>I. Sychugov, R. Juhasz, J. Valenta, and J. Linnros, *Phys. Rev. Lett.* **94**, 087405 (2005).

<sup>6</sup>H. J. Krenner, E. C. Clark, T. Nakaoka, M. Bichler, C. Scheurer, G. Abstreiter, and J. J. Finley, *Phys. Rev. Lett.* **97**, 076403 (2006).

<sup>7</sup>P. Ester, S. Stuffer, S. M. de Vasconcellos, M. Bichler, and A. Zrenner, *Phys. Status Solidi C* **3**, 3722 (2006).

<sup>8</sup>J. G. Keizer, J. Bocquel, P. M. Koenraad, T. Mano, T. Noda, and K. Sakoda, *Appl. Phys. Lett.* **96**, 062101 (2010).

<sup>9</sup>J. H. Blokland, M. Bozkurt, J. M. Ulloa, D. Reuter, P. M. Koenraad, P. C. M. Christianen, and J. C. Maan, *Appl. Phys. Lett.* **94**, 023107 (2009).

<sup>10</sup>V. Mlinar, M. Bozkurt, J. M. Ulloa, M. Ediger, G. Bester, A. Badolato, P. M. Koenraad, R. J. Warburton, and A. Zunger, *Phys. Rev. B* **80**, 165425 (2009).

<sup>11</sup>D. P. Kumah, S. Shusterman, Y. Paltiel, Y. Yacoby, and R. Clarke, *Nat. Nano.* **4**, 835 (2009); D. P. Kumah, J. H. Wu, N. S. Hussein, V. D. Dasika, R. S. Goldman, Y. Yacoby, and R. Clarke, *Appl. Phys. Lett.* **98**, 021903 (2011). A more accurate structure profile of epitaxy QDs can be indirectly obtained from a full three-dimensional electron density map measured by a coherent Bragg rod analysis (COBRA) method.

<sup>12</sup>F. Hatami, W. T. Masselink, L. Schrottke, J. W. Tomm, V. Talalaev, C. Kristukat, and A. R. Goñi, *Phys. Rev. B* **67**, 085306 (2003).

<sup>13</sup>N. Koguchi, S. Takahashi, and T. Chikyow, *J. Cryst. Growth* **111**, 688 (1991); N. Koguchi and K. Ishige, *Jpn. J. Appl. Phys., Part I* **32**, 2052 (1993).

<sup>14</sup>K. Kowalik, O. Krebs, A. Lemaître, J. A. Gaj, and P. Voisin, *Phys. Rev. B* **77**, 161305(R) (2008).

<sup>15</sup>M. Jo, T. Mano, and K. Sakoda, *J. Appl. Phys.* **108**, 083505 (2010).

<sup>16</sup>K. Kuroda, T. Kuroda, K. Sakoda, K. Watanabe, N. Koguchi, and G. Kido, *Appl. Phys. Lett.* **88**, 124101 (2006).

<sup>17</sup>M. Abbarchi, C. A. Mastrandrea, T. Kuroda, T. Mano, K. Sakoda, N. Koguchi, S. Sanguinetti, A. Vinattieri, and M. Gurioli, *Phys. Rev. B* **78**, 125321 (2008).

<sup>18</sup>T. Belhadji, *Appl. Phys. Lett.* **97**, 051111 (2010).

<sup>19</sup>T. Mano, M. Abbarchi, T. Kuroda, C. A. Mastrandrea, A. Vinattieri, S. Sanguinetti, K. Sakoda, and M. Gurioli, *Nanotechnology* **20**, 395601 (2009).

<sup>20</sup>M. Abbarchi, T. Kuroda, C. Mastrandrea, A. Vinattieri, S. Sanguinetti, T. Mano, K. Sakoda, and M. Gurioli, *Physica E* **42**, 881 (2010).

<sup>21</sup>J. W. Luo, G. Bester, and A. Zunger, *Phys. Rev. B* **79**, 125329 (2009).

<sup>22</sup>A. Franceschetti, H. Fu, L. W. Wang, and A. Zunger, *Phys. Rev. B* **60**, 1819 (1999).

<sup>23</sup>We are grateful to P. M. Koenraad and M. Takaaki for clarifying to us now that the dots used in XSTM were different than those used for PL measurements.

<sup>24</sup>J. G. Keizer, M. Jo, T. Mano, T. Noda, K. Sakoda, and P. M. Koenraad, *Appl. Phys. Lett.* **98**, 193112 (2011).

<sup>25</sup>V. Mlinar and A. Zunger, *Phys. Rev. B* **80**, 205311 (2009).

<sup>26</sup>V. Mlinar and A. Zunger, *Phys. Rev. B* **80**, 035328 (2009).

<sup>27</sup>E. L. Ivchenko, *Phys. Status Solidi A* **164**, 487 (1997).

<sup>28</sup>R. Singh and G. Bester, *Phys. Rev. Lett.* **104**, 196803 (2010).

<sup>29</sup>J. W. Luo, A. Franceschetti, and A. Zunger, *Phys. Rev. B* **79**, 201301(R) (2009).

<sup>30</sup>J. D. Plumhof, V. K2