Electrically-controlled two-dimensional gratings based on layers undulations in cholesteric liquid crystals

Bohdan Senyuk^{1,*}, Ivan Smalyukh², Oleg Lavrentovich^{1,2,}
¹Chemical Physics Interdisciplinary Program and ²Liquid Crystal Institute,
Kent State University, Kent, OH, USA 44242-0001

ABSTRACT

Liquid Crystals IX, edited by Iam-Choon Khoo, Proceedings of SPIE Vol. 5936 (SPIE, Bellingham, WA, 2005) \cdot 0277-786X/05/\$15 \cdot doi: 10.1117/12.615976

Proc. of SPIE Vol. 5936 59360W-1

2. EXPERIMENTAL

2.1. Cell preparation and materials

The LC cells for 2D cholesteric gratings (Fig. 1) were constructed from glass substrates coated with transparent indiumtin-oxide (ITO) patterned electrodes. The unidirectionally buffed thin layers of polyimide PI2555 (HD MicroSystem) were used to set the easy axis for LC molecules at the confining glass plates; we used different rubbing

Proc. of SPIE Vol. 5936 59360W-2

The LC cells (Fig. 1) were filled with cholesteric mixtures (Table 1) in the isotr

of oily streaks (Fig. 3). In order to prevent nucleation at the cell edges, the transparent electrodes were patterned (Fig. 1) with the working area being 25 mm^2 . This allows us to avoid the oily streaks by eliminating the nucleation sites such as surface irregularities, mechanical impurities, and strong layers distortions at the cell edges. The obtained 2D undulations were stable in time without being spoiled by the oily streaks.

$$m = L_g \sin ag{1}$$

where m is the diffraction order, and $L_{\rm g}$

period $L_u/2$ and the beams "s" (Fig. 4(b)) are caused by modulation of refractive index close to surfaces with period L_u . When d/p is large, the contribution from the regions close to the surfaces is negligib	

- 12. Anup K. Ghosh, Yoichi Takanishi, Ken Ishikawa, and Hideo Takezoe, Youshiyuki Ono, Joji Kawamura, "Electrically controllable polarization-dependent phase grating from photocurable liquid crystals," *J. Appl. Phys.* **95**(9), 5241-5243, 2004.
- 13. B. I. Senyuk, I. I. Smalyukh, O. D. Lavrentovich, "Switchable two-dimensional gratings based on field-induced layer undulations in cholesteric liquid crystals," *Opt. Lett.* **30**(4), 349-351, 2005.
- 14. I. I. Smalyukh, S. V. Shiyanovskii, and O. D. Lavrentovich, "Three-dimensional imaging of orientational order by fluorescence confocal polarizing microscopy," *Chem. Phys. Lett.* **336**, 88-96, 2001.
- 15. T. S. Lubensky, "Hydrodynamics in cholesteric liquid crystals," *Phys. Rev. A* 6, 452-470, 1972.
- 16. P.G. de Gennes and J. P002.O,