

Multimodal nonlinear optical polarizing microscopy of long-range molecular order in liquid crystals


Taewoo Lee, Rahul P. Trivedi, and Ivan I. Smalyukh*

Department of Physics, University of California, Santa Barbara, California 93106, USA
 *Corresponding author: smalyukh@ucsb.edu

Received August 26, 2010; accepted September 9, 2010;
 posted September 27, 2010 (Doc. ID 134049); published October 12, 2010

We demonstrate orientation-sensitive multimodal nonlinear optical polarizing microscopy capable of probing orientational, polar, and biaxial features of mesomorphic ordering in soft matter. This technique achieves simultaneous imaging in broadband coherent anti-Stokes Raman scattering, multiphoton excitation fluorescence, and multiharmonic generation polarizing microscopy modes and is based on the use of a single femtosecond laser and a photonic crystal fiber as sources of the probing light. We show the viability of this technique for mapping of three-dimensional patterns of molecular orientations and show that images obtained in different microscopy modes are consistent with each other. © 2010 Optical Society of America

OCIS codes: 110.1650, 160.3710, 180.4315, 180.6900.



C* (F \1 015-100), A (8CB, 4-4-4-4)
) 5CB (4-4-4-4)
EM C , F A CA M LC CN
(2236 -1) 8CB 5CB
F 2 HG
(3 1) 1050 C*

3 F, P - P P P P P P CA - M,